EYEM = * P31 92.006078/01 *SU 1627.112.A

Amblyopia-organic optic tract pathology differential diagnosis rhythmic stimulation with red light flashes with 30.35 hz frequency

EYE MICROSURGERY 22.06.88-SU-445846

S05 (15.02.91) A61b-03

22.06.88 as 445846 (1462AK)

The method involves recording induced visual potentials in response

to rhythmic light stimulation.

Rhythmic stimulation is performed with series of flashes of red light with a frequency of 30-35 Hz and a duration of 0.6-1.0s and if the induced potential has a frequency equal to that of the stimulating signal and an amplitude corresp. to the norm, then a diagnosis of amblyopia is made, whereas if there is no reproduction of the rhythm of the stimulating signal and if the amplitude of the induced potential is reduced, then organic pathology of the optic tract is diagnosed.

USE/ADVANTAGE - For differential diagnosis of amblyopia and organic pathology of the optic tract. More accurate method is

obtained. Bul.6/15.2.91. (2pp Dwg.No.0/0)

N92-004628

© 1992 DERWENT PUBLICATIONS LTD.

128, Theobalds Road, London WC1X 8RP, England
US Office: Derwent Inc., 1313 Dolley Madison Boulevard,
Suite 303, McLean, VA22101, USA
Unauthorised copying of this abstract not permitted.

BEST AVAILABLE COPY